

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

to make 100. By trial we find that $\frac{3}{6} \times 8 + \frac{6}{6} \times 7 = 100$. Hence, multiplying the column of 8 by $\frac{3}{6}$ and the column of 7 by $\frac{6}{6}$ and adding the results horizontally, we get 3 lambs, 68 sheep, and 29 hogs and calves. In like manner, by multiplying the column of 8 by $\frac{1}{6}$ and the column of 7 by $\frac{6}{6}$, we get 10 lambs, 60 sheep, and 30 hogs and calves. By multiplying by $\frac{1}{6}$ and $\frac{5}{6}$ respectively, we get 17 lambs, 52 sheep, and 31 hogs and calves. The remaining six answers may now be easily written down without further trial. Since, in the first result there are 29 hogs and calves, we may have 1 hog and 28 calves, and so on. In all we may have 28 different results. In like manner in the second result, we may have 29 different results. In like manner 30 in third, 31 in fourth, 32 in the fifth, 33 in the sixth, 34 in the seventh, 35 in the eighth and 36 in the ninth. Hence, in all, we have 28 + 29 + 30 + 31 + 32 + 33 + 34 + 35 + 36 = 288 different results.

For a fuller treatment of this class of problems see my Mathematical Solution Book.

Correct solutions to this problem were received from H. J. Geartner Martin Spinx, John McDowell, P_{\star} S. Berg. P. C. Cullen, and M. A. Gruber. Mr. Gruber also gave a solution by Algebra,


32. Proposed by P. C. CULLEN, Meade, Nebraska.

 Λ horse is tied to a corner of a building 40 feet square, by a rope 110 feet long. Over how much land can be graze?

Solution by MARTIN SPINX, Wilmington, Ohio; Professor A. J. LILLY, Algona, Iowa; and Professor H. J. GEARTNER, Wilmington, Ohio

Let ABCD represent the barn, side AB=40 feet; BE represent the rope, length 110 feet; and CG=70 feet. Then the area of $FBEKF=\frac{3}{4}\pi\times110^2$ = 28510.02 sq. feet. The area of the two equal quadrants AFGH and $CLGE=\frac{1}{2}\pi70^2=7696.84$ sq. feet. Now $AC=40\sqrt{2}$ feet; $PG=\sqrt{(GC^2-PC^2)}$

= $\sqrt{(70^2-800)}=10_1$ 41 feet; and $DG=10_1/41-20_1/2=10[\sqrt{41-2\sqrt{2}}]$ feet, ... Area of square $ODIG=GD^2\div2=\frac{1}{3}10[\sqrt{41-2\sqrt{2}}]^{\frac{1}{3}}$ $\div2=638.922932$ sq. ft. The height of the segment GNL is LO=LD-OD=30 feet $-\frac{1}{2}\sqrt{2}$ $GD=[30-(5_1/82-20)]$ feet $-\frac{5}{3}[10-\sqrt{82}]$ feet and the base is $GN=2GO=\sqrt{2}GD$ $10[\sqrt{82}-4]$ feet. Hence, according to the rule, p. 389, Ray's Higher Arithmetic, the area of the segments GNL and GMH is $2(LO^3\div2GN+\frac{2}{3}LO\times GN)=2\frac{1}{3}[5(10-\sqrt{82})]^3\div20[\sqrt{82}-4]+\frac{2}{3}\times5(10-\sqrt{82})\times[10_1/82-4]+\frac{2}{3}=320.443888$ sq. ft. and the



area of the half segments=160.221944 sq. ft. Hence, the area over which the horse grazes= $\frac{3}{4}FBEKF+FAH+LCE-(LOG+ODIG+GIH)=\frac{3}{4}\pi110^2+\frac{1}{2}\pi70^2-(160.221944+638.922932)=35407.72514$ sq. ft.

This problem was solved in different ways and with different results by $P,\,S,\,BERG,\,John\,McDowell,\,G,\,B,\,M,\,Zerr,\,and\,D,\,G,\,Durrance,$

[For a demonstration of the rule referred to above, see my Mathematical Solution Book, pp. 201 and 202.—Ed.]